bs-14995R

[Primary Antibody]

TENT5A Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –		400-901-9800
Host: Rabbit Clonality: Polyclonal	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
GenelD: 55603	SWISS: Q96IP4	ICC/IF (1:100-500) ELISA (1:5000-10000)
Target: TENT5A Immunogen: KLH conjugated synthetic peptide derived from human TENT5A: 201-300/442.		Reactivity: (predicted: Human, Mouse, Rat, Cow, Dog)
Purification: affinity purified by	Protein A	
Concentration: 1mg/ml		Predicted
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted 50 kDa MW.: ^{50 kDa} Subcellular Location: Cytoplasm ,Nucleus
contains around 1, sequence. Deletior associated with ea presence of a cance is associated with of when mutated, pre porphyria. Notably Parkinson's diseas histocompatibility components of the to rheumatic disea syndrome, 21-hydr disease are also as disorder susceptib chromosome 6. Th	% of the human genome, chromosome 6 200 genes within 170 million base pairs of a of a portion of the q arm of chromosome 6 is rly onset intestinal cancer suggesting the er susceptibility locus. Porphyria cutanea tarda chromosome 6 through the HFE gene which, disposes an individual to developing this , the PARK2 gene, which is associated with e, and the genes encoding the major complex proteins, which are key molecular immune system and determine predisposition ses, are also located on chromosome 6. Stickler oxylase deficiency and maple syrup urine sociated with genes on chromosome 6. A bipolar ility locus has been identified on the q arm of e FAM46A gene product has been provisionally A pending further characterization.	