


**bs-21389R****[ Primary Antibody ]****Utrophin Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 7402**SWISS:** P46939**Target:** Utrophin**Immunogen:** KLH conjugated synthetic peptide derived from human Utrophin: 565-665/3433.**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** Dystrophin and utrophin are related structural, Actin-binding proteins that are involved in anchoring the cytoskeleton to the plasma membrane. Dystrophin is the protein product of the Duchenne/Becker muscular dystrophy gene. Dystrophin expression is found in muscle and brain tissues, where it is localized to the inner surface of the plasma membrane. It has been speculated that alternative splicing of the carboxy terminus allows dystrophin to interact with a variety of proteins. Research has shown that the loss of dystrophin-associated proteins in Duchenne afflicted muscle is due to the absence of dystrophin rather than to muscle degradation and that the lack of dystrophin results in the loss of linkage between the cytoskeleton and the extracellular matrix. Evidence suggests that the upregulation of utrophin can reduce the dystrophic pathology.**Applications:** IHC-P (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Mouse (predicted: Human, Rabbit, Sheep, Cow, Chicken, Dog)**Predicted MW.:** 394 kDa**Subcellular Location:** Cell membrane ,Cytoplasm**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Utrophin) Polyclonal Antibody, Unconjugated (bs-21389R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Utrophin) Polyclonal Antibody, Unconjugated (bs-21389R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.