### bs-3436R

## [ Primary Antibody ]

# BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

IHC-F (1:100-500)

Chicken, Horse)

**Subcellular Location:** Cell membrane ,Cytoplasm

Mouse, Rabbit, Pig, Cow,

Reactivity: Rat (predicted: Human,

67 kDa

Applications: IHC-P (1:100-500)

Predicted

MW.:

## phospho-TAK1 (Thr184) Rabbit pAb

- DATASHEET -

**Host:** Rabbit **Isotype:** IgG

Clonality: Polyclonal

**GenelD:** 6885 **SWISS:** 043318

Target: TAK1 (Thr184)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

TAK1 around the phosphorylation site of Thr184: IQ(p-T)HM.

**Purification:** affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated


freeze/thaw cycles.

**Background:** The protein encoded by this gene is a member of the

serine/threonine protein kinase family. This kinase mediates the signaling transduction induced by TGF beta and morphogenetic protein (BMP), and controls a variety of cell functions including transcription regulation and apoptosis. In response to IL-1, this protein forms a kinase complex including TRAF6, MAP3K7P1/TAB1 and MAP3K7P2/TAB2; this complex is required for the activation of nuclear factor kappa B. This kinase can also activate MAPK8/JNK, MAP2K4/MKK4, and thus plays a role in the cell response to environmental stresses. Four alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided

by RefSeq, Jul 2008]

VALIDATION IMAGES



Tissue/cell: rat brain tissue; 4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-Phospho-TAK1(Thr184) Polyclonal Antibody, Unconjugated(bs-3436R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

#### - SELECTED CITATIONS -

• [IF=3.36] Johnson Chacko L et al. Early appearance of key transcription factors influence the spatiotemporal development of the human inner ear. Cell Tissue Res. 2019 Dec 2. IF; Human fetuses. 31788757