

**bs-1208R****[ Primary Antibody ]****ADAMTS1 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>WB</b> (1:500-2000)
<b>Clonality:</b> Polyclonal		<b>IHC-P</b> (1:100-500)
<b>GeneID:</b> 11504	<b>SWISS:</b> P97857	<b>IHC-F</b> (1:100-500)
<b>Target:</b> ADAMTS1		<b>IF</b> (1:100-500)
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from mouse ADAMTS1: 851-967/967.		<b>Reactivity:</b> Human (predicted: Mouse, Rat, Pig, Sheep, Cow, Chicken, Horse)
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		<b>Predicted MW.:</b> 106 kDa
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		<b>Subcellular Location:</b> Secreted ,Extracellular matrix
<b>Background:</b> ADAMTS1 is a metalloproteinase of the ADAM (A Disintegrin And Metalloproteinase) family containing disintegrin-like domains. ADAMTS1, also known as METH1, was first described as a protein elevated in invasive mouse tumors. Initial findings indicated a role for ADAMTS1 in tumor progression, since the protein was preferentially expressed in more invasive tumor cell lines. ADAMTS1 is necessary for normal growth, fertility, and organ morphology and function.		

**— VALIDATION IMAGES —**

Sample: Lane 1: Human Hela cell Lysates Lane 2: Human 293T cell Lysates Lane 3: Human U87-MG cell Lysates Lane 4: Human U251 cell Lysates  
Primary: Anti-ADAMTS1 (bs-1208R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 106kDa Observed band size: 80kDa



Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-ADAMTS1 Polyclonal Antibody, Unconjugated(bs-1208R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

**— SELECTED CITATIONS —**

- **[IF=3]** Hiroe Toba. et al.Suppressing SPARC gene with siRNA exerts therapeutic effects and inhibits MMP-2/9 and ADAMTS1 overexpression in a murine model of ischemia/reperfusion-induced acute kidney injury.Journal of Pharmacological Sciences. IHC ;Mouse. 10.1016/j.jphs.2025.03.010