

**bs-13715R****[ Primary Antibody ]****CDH12 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<p><b>Host:</b> Rabbit</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>GeneID:</b> 1010</p> <p><b>Target:</b> CDH12</p> <p><b>Immunogen:</b> KLH conjugated synthetic peptide derived from human CDH12: 291-390/794. &lt; Extracellular &gt;</p> <p><b>Purification:</b> affinity purified by Protein A</p> <p><b>Concentration:</b> 1mg/ml</p> <p><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> The cadherins are a family of Ca<sup>++</sup>-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morpho-genesis (1-4). Cadherins each contain a large extracellular domain at the amino terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity (5,6). The relatively short carboxy terminal, intracellular domain interacts with a variety of cytoplasmic proteins, including catenin b, to regulate cadherin function (7). BR-cadherin (for brain-cadherin, also designated cadherin-12 or N-cadherin 2) is expressed specifically in neurons in the central nervous system and is thought to be involved in neuronal development (8).</p>	<p><b>Applications:</b> <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500)</p> <p><b>Reactivity:</b> Rat (predicted: Human, Mouse, Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)</p> <p><b>Predicted MW.:</b> 82 kDa</p> <p><b>Subcellular Location:</b> Cell membrane</p>
---	---

**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CDH12) Polyclonal Antibody, Unconjugated (bs-13715R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.