

**bs-0433R****[ Primary Antibody ]****ADAMTS7 Rabbit pAb****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500)
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 11173	<b>SWISS:</b> Q6P7J9	
<b>Target:</b> ADAMTS7		<b>Reactivity:</b> Human, Rat (predicted: Mouse, Cow)
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human ADAMTS7: 1601-1686/1686.		
<b>Purification:</b> affinity purified by Protein A		<b>Predicted MW.:</b> 158 kDa
<b>Concentration:</b> 1mg/ml		<b>Subcellular Location:</b> Secreted ,Extracellular matrix ,Cell membrane
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Metalloprotease that may play a role in the degradation of COMP. Binds 1 zinc ion per subunit. pH dependence: Optimum pH is between 7.5 and 9.5. [Subcellular location] Secreted, extracellular space, extracellular matrix (By similarity). Note=Also found associated with the external cell surface[Tissue specificity] Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Detected in meniscus, bone, tendon, cartilage, synovium, fat and ligaments. Up-regulated in articular cartilage and synovium from arthritis patients.		

**— VALIDATION IMAGES —**

Tissue/cell: human placenta tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-ADAMTS7 Polyclonal Antibody, Unconjugated(bs-0433R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-ADAMTS7 Polyclonal Antibody, Unconjugated(bs-0433R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining