[Primary Antibody]

phospho-ASK1 (Thr845) Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

| – DATASHEET – | | 400-901-9800 |
|---|--|--|
| Host: Rabbit | lsotype: lgG | Applications: WB (1:500-2000) |
| Clonality: Polyclonal | | Flow-Cyt (1µg/test) |
| GenelD: 4217 | SWISS: Q99683 | Reactivity: Human, Mouse, Rat |
| Target: ASK1 (Thr845) | | |
| Immunogen: KLH conjugated S ASK1 around the p | nthesised phosphopeptide derived from human hosphorylation site of Thr845: TE(p-T)FT. | Predicted |
| Purification: affinity purified by Protein A | | MW.: ^{155 kDa} Subcellular Location: ^{Cytoplasm} |
| Concentration: 1mg/ml | | |
| Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | | |
| Background: Mitogen-activated include MAPK or e kinase (MKK or ME MAPKK kinase/ME protein kinase, MA The kinases of the homologs exist in MAPKKK5 contain subdomains. Nort transcript is abund The MAPKKK5 pro SERK1, MAPKK4) i (JNK)/stress-activ expression in COS | protein kinase (MAPK) signaling cascades xtracellular signal-regulated kinase (ERK), MAPK K), and MAPK kinase kinase (MAPKKK or MEKK). KK phosphorylates and activates its downstream PK kinase/MEK, which in turn activates MAPK. se signaling cascades are highly conserved, and yeast, Drosophila, and mammalian cells. s 1,374 amino acids with all 11 kinase hern blot analysis shows that MAPKKK5 dantly expressed in human heart and pancreas. tein phosphorylates and activates MKK4 (aliases n vitro, and activates c-Jun N-terminal kinase ated protein kinase (SAPK) during transient and 293 cells; MAPKKK5 does not activate | |

- VALIDATION IMAGES

MÁPK/ERK.



Sample: Lane 8: Cerebrum (Rat) Lysate at 40 ug Lane 9: Stomach (Rat) Lysate at 40 ug Primary: Anti-phospho-ASK1 (Thr845) (bs-3031R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 154 kD Observed band size: 154 kD Sample: Lane1: Brain(Rat) Lysate at 30 ug Lane2: Heart(Rat) Lysate at 30 ug Primary: Antiphospho-ASK1(Thr845) (bs-3031R) at 1:200 dilution; Secondary: HRP conjugated Goat-Anti-Rabbit IgG(bs-0295G-HRP) at 1: 3000 dilution; Predicted band size : 155kD Observed band size :

155kD



Blank control:Mouse spleen. Primary Antibody (green line): Rabbit Anti-phospho-RelB (Ser551) antibody (bs-3031R) Dilution: $2\mu g\,/10^{\Lambda}6$ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF488 Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at-20°C. The cells were then incubated in 5%BSA to block nonspecific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control:U937. Primary Antibody (green line): Rabbit Anti-phospho-ASK1 (Thr845) antibody (bs-3031R) Dilution: 1ug/Test; Secondary Antibody : Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block nonspecific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=9.58] Xiaoli Guo. et al. ASK1 signaling regulates phase-specific glial interactions during neuroinflammation. P Natl Acad Sci Usa. 2022 Feb;119(6): IHC ;MOUSE. 35101972
- [IF=5.193] Zhou D et al. Inhibition of apoptosis signal-regulating kinase by paeoniflorin attenuates neuroinflammation and ameliorates neuropathic pain. J Neuroinflammation. 2019 Apr 11;16(1):83. WB ;Rat. 30975172
- [IF=5.1] Mengni He. et al. Exploring novel indazole derivatives as ASK1 inhibitors: Design, synthesis, biological evaluation and docking studies. BIOORG CHEM. 2024 Jun;147:107391 WB ;Human. 38677010
- [IF=5.014] Ying-Jung Hsu. et al. Protective Effect of Fenofibrate on Oxidative Stress-Induced Apoptosis in Retinal–Choroidal Vascular Endothelial Cells: Implication for Diabetic Retinopathy Treatment. Antioxidants-Basel. 2020 Aug;9(8):712 WB ;Monkey. 32764528
- [IF=3.569] Choi H et al. Apoptosis signal-regulating kinase 1 activation by Nox1-derived oxidants is required for TNFα receptor endocytosis. Am J Physiol Heart Circ Physiol. 2019 Mar 29. WB ;MOUSE. 30925081