bsm-34147M

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

ACTH Mouse mAb

- DATASHEET -

Host: Mouse **Isotype:** IgG1, Kappa

Clonality: Monoclonal CloneNo.: 7D13

GeneID: 5443 SWISS: P01189

Target: ACTH

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: Liquid in PBS containing 50% Glycerol, 0.5% BSA and 0.02%

Proclin300.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene encodes a polypeptide hormone precursor that

undergoes extensive, tissue-specific, post-translational processing via cleavage by subtilisin-like enzymes known as prohormone convertases. There are eight potential cleavage sites within the polypeptide precursor and, depending on tissue type and the available convertases, processing may yield as many as ten biologically active peptides involved in diverse cellular functions. The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation. These include several distinct melanotropins, lipotropins, and endorphins that are contained within the adrenocorticotrophin and beta-lipotropin peptides. Mutations in this gene have been associated with early onset obesity, adrenal insufficiency, and red hair pigmentation. Alternatively spliced transcript variants encoding the same protein have been described. [provided by

Applications: IHC-P (1:100-500)


IHC-F (1:100-500) IF (1:100-500)

Reactivity: Human

Subcellular Location: Secreted

VALIDATION IMAGES

RefSeq, Jul 2008].



Paraformaldehyde-fixed, paraffin embedded (Human pituitary adenoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (ACTH) Monoclonal Antibody, Unconjugated (bsm-34147M) at 1:100 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.