

bsm-34153M**[Primary Antibody]**

BioSS
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

NAPSIN A Mouse mAb**— DATASHEET —**

Host: Mouse	Isotype: IgG1, Kappa	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200) Reactivity: Human, Mouse, Rat Predicted MW.: 39 kDa Subcellular Location: Secreted
Clonality: Monoclonal	CloneNo.: 10A12	
Target: NAPSIN A		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: Liquid in PBS containing 50% Glycerol, 0.5% BSA and 0.02% Proclin300. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Napsin A is an aspartic proteinase that belongs to the peptidase A1 family and plays a role in pneumocyte surfactant processing. It is a 420-amino acid polypeptide consisting of a 24-residue signal peptide, a 40-amino acid propeptide, the mature enzyme of 336 amino acids, and a C-terminal extension of 18 residues. The mature Napsin A protein contains 3 predicted disulfide bonds, 3 potential N-linked oligosaccharide attachment sites, an RGD motif, a recognition motif for integrin binding, in the C terminus, immediately before a 4-amino acid insert that is unique to aspartic proteinases. Highest levels of Napsin A have been detected in adult lung (type II pneumocytes), fetal lung, and kidney tissues. Napsin A is also expressed at lower levels in adult spleen and at very low levels in peripheral blood leukocytes. Human napsin A shares 72.6% sequence identity with the mouse homolog.		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (NAPSIN A) Monoclonal Antibody, Unconjugated (bsm-34153M) at 1:100 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (NAPSIN A) Monoclonal Antibody, Unconjugated (bsm-34153M) at 1:100 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.