## bs-18782R

## [ Primary Antibody ]

## Mel18 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		
Host: Rabbit Clonality: Polyclonal	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500)
GenelD: 7703 Target: Mel18	SWISS: P35227	IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000)
Immunogen: KLH conjugated synthetic peptide derived from human Mel18: 281-344/344.		Rat, Rabbit, Pig, Sheep,
<b>Purification:</b> affinity purified by <b>Concentration:</b> 1mg/ml	Protein A	Cow, Dog, Horse, Chimpanzee, Macaque Monkey)
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted <sub>38 kDa</sub> MW.: <sup>38 kDa</sup>
<b>Background:</b> The protein encoded by this gene contains a RING finger motif and is similar to the polycomb group (PcG) gene products. PcG gene products form complexes via protein-protein interaction and maintain the transcription repression of genes involved in embryogenesis, cell cycles, and tumorigenesis. This protein was shown to act as a negative regulator of transcription and has tumor suppressor activity. The expression of this gene was detected in various tumor cells, but is limited in neural organs in normal tissues. Knockout studies in mice suggested that this protein may negatively regulate the expression of different cytokines, chemokines, and chemokine receptors, and thus plays an important role in lymphocyte differentiation and migration, as well as in immune responses. [provided by RefSeq, Jul 2008]		gene d was is is is in is in is in is in is