

**bs-55192R**

**[ Primary Antibody ]**

## SAV1 Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

### — DATASHEET —

<p><b>Host:</b> Rabbit</p> <p><b>Clonality:</b> Polyclonal</p> <p><b>GeneID:</b> 60485</p> <p><b>Target:</b> SAV1</p> <p><b>Immunogen:</b> Recombinant human SAV1 protein: 199-383/383.</p> <p><b>Concentration:</b> 1mg/ml</p> <p><b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p><b>Background:</b> WW domain-containing proteins are found in all eukaryotes and play an important role in the regulation of a wide variety of cellular functions such as protein degradation, transcription, and RNA splicing. This gene encodes a protein with two WW domains, a SARA domain, and a coiled-coil region and is ubiquitously expressed in adult tissues. This protein binds to MST1 (mammalian sterile 20-like kinase 1) and promotes MST1-induced apoptosis. It has also been shown to bind to HAX1 (hematopoietic cell-specific protein 1 (HS1)-associated protein X-1) and to attenuate the anti-apoptotic effects of HAX1. Studies in human and mouse suggest this gene acts as a tumor suppressor. [provided by RefSeq, Aug 2012]</p>	<p><b>Isotype:</b> IgG</p> <p><b>SWISS:</b> Q9H4B6</p>	<p><b>Applications:</b> <b>WB</b> (1:1000 - 1:4000) <b>ELISA</b> (1:5000-10000)</p> <p><b>Reactivity:</b> Human, Mouse</p> <p><b>Predicted MW.:</b> 45 kDa</p> <p><b>Subcellular Location:</b> Cytoplasm ,Nucleus</p>
--	--	---