bs-0253R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Estrogen receptor alpha Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 2099 SWISS: P03372

Target: Estrogen receptor alpha

Immunogen: KLH conjugated synthetic peptide derived from human Estrogen

Receptor alpha: 241-300/595.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Estrogen and progesterone receptor are members of a family of transcription factors that are regulated by the binding of their cognate ligands. The interaction of hormone-bound estrogen receptors with estrogen responsive elements(EREs) alters transcription of ERE-containing genes. The carboxy terminal region of the estrgen receptor contains the ligand binding domain, the amino terminus serves as the transactivation domain, and the DNA binding domain is centrally located. Two forms of estrogen receptor have been identified, ER Alpha and ER Beta. ER Alpha and ER Beta have been shown to be differentially activated by various ligands. The biological response to progesterone is mediated by two distinct forms of the human progesterone receptor (hPR-A and hPR-B), which arise from alternative splicing. In most cells, hPR-B functions as a transcriptional activator of progesterone-responsive gene, whereas hPR-A function as a transcriptional inhibitor of all steroid hormone receptors.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500) ICC/IF (1:100-500) **ELISA** (1:5000-10000)

Reactivity: Human (predicted: Mouse,


Rat)

Predicted MW.: 66 kDa

Subcellular Cell membrane, Cytoplasm

Location: , Nucleus

VALIDATION IMAGES



4% Paraformaldehyde-fixed MCF-7 (H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (Estrogen receptor alpha) polyclonal Antibody, unconjugated (bs-0253R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-40295G-FITC) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.

— SELECTED CITATIONS –

• [IF=2.65] Zaki Asmaa Hussein. et al. Medicinal Mushroom Leucocalocybe mongolica Imai Extracts Improve Mammary Gland Differentiation in Lactating Rats via Regulating Protein Expression. EVID-BASED COMPL ALT. 2022;2022:5762847 IHC ;Rat. 35761899

- [IF=2.089] Huan He. et al. BDE-209 disturbed proliferation and differentiation of spermatogonia during mitotic process through estrogen receptor α. REPROD BIOL. 2023 Jun;23:100737 WB,IF;Rat. 36821943
- [IF=1.487] Mengmeng Yu. et al. Daidzein promotes milk synthesis and proliferation of mammary epithelial cells via the estrogen receptor α-dependent NFκB1 activation. 2020 May 13 WB,IF; Bovine. 32401613