
CYPA Rabbit pAb

Catalog Number: bs-5912R

Target Protein: CYPA

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), Flow-Cyt (1µg/Test)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Pig, Cow, Dog)

Predicted MW: 18 kDa

Subcellular Cytoplasm

Locations:

Entrez Gene: 5478

Swiss Prot: P62937

Source: KLH conjugated synthetic peptide derived from human Cyclophilin A.: 101-165/165.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The immunosuppressant cyclosporin A (CsA) forms a trimolecular complex with cyclophilin and calcineurins to inhibit calcineurin phosphatase activity (1). Cyclophilins are conserved, ubiquitous and abundant cytosolic peptidyl-prolyl cis-trans isomerases that accelerate the isomerization of XaaPro peptide bonds and the refolding of proteins (2,3). Human cyclophilin A (CyPA), an intracellular protein of 165 amino acids, is the target of the CsA and is encoded by a single unique gene conserved from yeast to humans (4,5). CyPA is known for its involvement in T cell differentiation and proliferation and is highly expressed in brain (6). CyPA is incorporated into the virion of the type 1 human immunodeficiency virus (HIV-1) via a direct interaction with the capsid domain of the viral Gag polyprotein and is crucial for efficient viral replication (7,8). Cyclophilin B (CyPB) is a member of the cyclophilin family with specific N- and C-terminal extensions. Unlike CyPA, CyPB has a signal sequence leading to its translocation in the endoplasmic reticulum. CyPB is secreted in biological fluids such as blood or milk and binds to a specific receptor present on the human lymphoblastic cell line Jurkat and on human peripheral blood lymphocytes (9,10).


VALIDATION IMAGES




Sample: Cerebrum (Rat) Lysate at 40 ug HeLa(Human) Cell Lysate at 30 ug Primary: Anti-CYPA (bs-5912R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 18 kD Observed band size: 17/18 kD




Sample: NIH/3T3(Mouse) Cell Lysate at 30 ug A431(Human) Cell Lysate at 30 ug Primary: Anti-CYPA (bs-5912R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 18 kD Observed band size: 18 kD




Sample: Lane 1: Mouse Thymus tissue lysates Lane 2: Mouse Lymph node tissue lysates Lane 3: Mouse Lung tissue lysates Lane 4: Mouse NIH/3T3 cell lysates Lane 5: Rat Thymus tissue lysates Lane 6: Rat Lymph node tissue lysates Lane 7: Rat Lung tissue lysates Lane 8: Human HeLa cell lysates Lane 9: Human PANC-1 cell lysates Lane 10: Human Molt-4 cell lysates Lane 11: Human HUVEC cell lysates Lane 12: Human THP-1 cell lysates Primary: Anti- CYPA (bs-5912R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 18 kDa Observed band size: 18 kDa



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-CYPA Polyclonal Antibody, Unconjugated(bs-5912R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-CYPA Polyclonal Antibody, Unconjugated(bs-5912R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control(blue): HeLa(fixed with 2% paraformaldehyde (10 min) , then permeabilized with 90% ice-cold methanol for 30 min on ice). Primary Antibody:Rabbit Anti- CYPA antibody(bs-5912R), Dilution: 1μg in 100 μL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions); Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.

PRODUCT SPECIFIC PUBLICATIONS

[IF=3.58] Kang, Taewook, et al. "Large-scale analysis of posttranslational modifications in the hippocampus of patients with Alzheimer' s disease using pI shift and label-free quantification without enrichment." Analytical and Bioanalytical Chemistry (2014): 1-14. WB ;
="Human" . 25120180