

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Calreticulin Rabbit pAb

Catalog Number: bs-5913R
Target Protein: Calreticulin

Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ELISA (1:5000-10000)

Reactivity: Human
Predicted MW: 44 kDa
Entrez Gene: 811
Swiss Prot: P27797

Source: KLH conjugated synthetic peptide derived from human Calreticulin: 101-200/417.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Calreticulin is a highly conserved chaperone protein which resides primarily in the


endoplasmic reticulum, and is involved in a variety of cellular processes, among them, cell adhesion. Additionally, it functions in protein folding quality control and calcium homeostasis. Calreticulin is also found in the nucleus, suggesting that it may have a role in transcription regulation. Systemic lupus erythematosus is associated with increased autoantibody titers against calreticulin. Recurrent mutations in calreticulin have been linked to various

neoplasms, including the myeloproliferative type.[provided by RefSeq, May 2020]

VALIDATION IMAGES



Sample: Lane 1: Human Jurkat cell lysates Lane 2: Human HepG2 cell lysates Lane 3: Human MCF-7 cell lysates Primary: Anti-Calreticulin (bs-5913R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 44 kDa Observed band size: 52 kDa



Paraformaldehyde-fixed, paraffin embedded Human Lung; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Calreticulin Polyclonal Antibody, Unconjugated (bs-5913R) at 1:200 overnight at 4° C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.




Paraformaldehyde-fixed, paraffin embedded Human Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Calreticulin Polyclonal Antibody, Unconjugated (bs-5913R) at 1:200 overnight at 4° C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Calreticulin Polyclonal Antibody, Unconjugated (bs-5913R) at 1:200 overnight at 4° C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Cerebrum; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Calreticulin Polyclonal Antibody, Unconjugated (bs-5913R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Calreticulin Polyclonal Antibody, Unconjugated (bs-5913R) at 1:200 overnight at 4° C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.

PRODUCT SPECIFIC PUBLICATIONS

[IF=29.4] Zhenyu Wang. et al. Laser-Activatable in Situ Vaccine Enhances Cancer-Immunity Cycle. ADV MATER. 2023 Nov;:2307193 FCM; MOUSE . 37951210

[IF=27.4] Qianyi Su. et al. Eradication of Large Tumors by Nanoscale Drug Self-Assembly. ADV MATER. 2024 Oct;:2410536 FC; MOUSe. 39420689

[IF=18.9] Ze Wang. et al. Turning foes to friends: Advanced "in situ nanovaccine" with dual immunoregulation for enhanced immunotherapy of metastatic triple-negative breast cancer. BIOACT MATER. 2024 Sep;39:612 FCM; MOUSE . 38883315

[IF=18.5] Baohua Zhang. et al. Self-Oxygenated Biomimetic Nanozyme for Tumor Catalytic Immunotherapy. ADV FUNCT MATER. 2024 Oct;:2411103 FC; MOUSE . 10.1002/adfm.202411103

[IF=17.694] Fu Shunli. et al. Temperature sensitive liposome based cancer nanomedicine enables tumour lymph node immune microenvironment remodelling. NAT COMMUN. 2023 Apr;14(1):1-17 IHC; Mouse. 37076492