bs-2215R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

phosphos-PCNA (Tyr211) Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 5111 SWISS: P12004

Target: phosphos-PCNA (Tyr211)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

PCNA around the phosphorylation site of Tyr211: LR(p-Y)LN.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Proliferating cell nuclear antigen (PCNA) is a 28kDa nuclear protein

associated with the cell cycle, a nuclear protein vital for cellular DNA synthesis. Proliferating cell nuclear antigen was originally identified by immunofluorescence as a nuclear protein whose appearance correlated with the proliferate state of the cell. PCNA is required for replication of DNA in vitro and has been identified as the auxiliary protein (cofactor) for DNA polymerase delta. The anti-PCNA antibodies react with the nuclei of proliferating cells. PCNA is essential for cellular DNA synthesis and is also required for the in vitro replication of simian virus 40 (SV40) DNA where it acts to coordinate leading and lagging strand synthesis at the replication fork. The PCNA protein may fulfil several separate roles in the cell nucleus associated with changes in its antigenic structure.

Applications: Flow-Cyt (1µg /test)

ICC/IF (1:100-500)

Reactivity: Human (predicted: Mouse,

Rat, Rabbit, Sheep, Cow,


Chicken)

Predicted 29 kDa


MW.:

Subcellular Nucleus

VALIDATION IMAGES



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (phosphos-PCNA (Tyr211)) polyclonal Antibody, Unconjugated (bs-2215R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control (blue line): U251 (fixed with 2% paraformaldehyde (10 min)and then permeabilized with 0.1% PBS-Tween for 20 min at room temperature). Primary Antibody (green line): Rabbit Anti-PCNA antibody (bs-2215R), Dilution: 1µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG. Secondary Antibody (white blue line): Goat antirabbit IgG-PE, Dilution: 1µg /test.