

bs-8394R**[Primary Antibody]****Fbxw7 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200) Flow-Cyt (1ug/Test) Reactivity: Human, Mouse, Rat (predicted: Pig, Cow, Dog, Horse) Predicted MW.: 78 kDa Subcellular Location: Cytoplasm ,Nucleus
Clonality: Polyclonal		
GeneID: 55294	SWISS: Q969H0	
Target: Fbxw7		
Immunogen: KLH conjugated synthetic peptide derived from human Fbxw7/CDC4: 501-600/707.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Fbw7 is a member of the F box protein family which are characterized by an approximately 40 amino acid motif, the F box. The F box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F box), which function in phosphorylation-dependent ubiquitination. The F box proteins are divided into 3 classes: Fbws containing WD40 domains, Fbls containing leucine rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. Fbw7 belongs to the Fbws class; in addition to an F box, this protein contains 7 tandem WD40 repeats. It binds directly to cyclin E and probably targets cyclin E for ubiquitin mediated degradation. Mutations of this gene are detected in ovarian and breast cancer cell lines. Alternative splicing of this gene generates 2 transcript variants diverging at the 5' termini.		

— VALIDATION IMAGES —


Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Fbxw7) Polyclonal Antibody, Unconjugated (bs-8394R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Fbxw7) Polyclonal Antibody, Unconjugated (bs-8394R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control (black line) :MCF-7. Primary Antibody (green line): Rabbit Anti-Fbxw7 antibody (bs-8394R) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control (black line) :MCF-7. Primary Antibody (green line): Rabbit Anti-Fbxw7 antibody (bs-8394R) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=8.74]** Chen et al. STAT1 inhibits human hepatocellular carcinoma cell growth through induction of p53 and Fbxw7. (2015) Cancer.Cell.Int. 15:111 WB ;Human. 26617467
- **[IF=7]** Chan Dedrick Kok Hong. et al. Biallelic FBXW7 knockout induces AKAP8-mediated DNA damage in neighbouring wildtype cells. CELL DEATH DISCOV. 2023 Jun;9(1):1-15 WB ;Human. 37386001
- **[IF=2.28]** Wang, Haihe, et al. "RBP-J-interacting and tubulin-associated protein induces apoptosis and cell cycle arrest in human hepatocellular carcinoma by activating the p53-Fbxw7 pathway." Biochemical and Biophysical Research Communications (2014). WB ;="Human". 25445601