

bs-15496R**[Primary Antibody]****ALDH3A1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 218	SWISS: P30838	IHC-F (1:100-500)
Target: ALDH3A1		IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide derived from human ALDH3A1: 151-250/453.		
Purification: affinity purified by Protein A		Reactivity: Mouse, Rat (predicted: Human, Pig, Horse)
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 50 kDa
Background: Aldehyde dehydrogenases oxidize various aldehydes to the corresponding acids. They are involved in the detoxification of alcohol-derived acetaldehyde and in the metabolism of corticosteroids, biogenic amines, neurotransmitters, and lipid peroxidation. The enzyme encoded by this gene forms a cytoplasmic homodimer that preferentially oxidizes aromatic and medium-chain (6 carbons or more) saturated and unsaturated aldehyde substrates. It is thought to promote resistance to UV and 4-hydroxy-2-nonenal-induced oxidative damage in the cornea. The gene is located within the Smith-Magenis syndrome region on chromosome 17. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Sep 2008].		Subcellular Location: Cytoplasm

— VALIDATION IMAGES —

Sample: Lung (Mouse) Lysate at 40 ug Primary: Anti-ALDH3A1 (bs-15496R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 50 kD
Observed band size: 50 kD



Sample: Stomach (Mouse) Lysate at 40 ug Primary: Anti-ALDH3A1 (bs-15496R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 50 kD Observed band size: 52 kD



Paraformaldehyde-fixed, paraffin embedded (rat stomach tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ALDH3A1) Polyclonal Antibody, Unconjugated (bs-15496R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

— SELECTED CITATIONS —

- **[IF=6.4]** Martin Schicht. et al. Ocular surface changes in mice with streptozotocin-induced diabetes and diabetic polyneuropathy. OCUL SURF. 2023 Dec;; IHC ;Mouse. 38141818