

bsm-33301M**[Primary Antibody]****phospho-STAT3 (Tyr705) Mouse mAb****Bioss**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


— DATASHEET —**Host:** Mouse**Isotype:** IgG**Clonality:** Monoclonal**CloneNo.:** 3F10**GeneID:** 6774**SWISS:** P40763**Target:** STAT3 (Tyr705)**Immunogen:** KLH conjugated Synthesised phosphopeptide derived from human STAT3 around the phosphorylation site of Tyr705: AP(p-Y)LK.**Purification:** affinity purified by Protein G**Concentration:** 1mg/ml

Storage: Size : 50ul/100ul/200ul
0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Size : 200ug (PBS only)
0.01M PBS
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. Mutations in this gene are associated with infantile-onset multisystem autoimmune disease and hyper-immunoglobulin E syndrome. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Sep 2015]

Applications: IHC-P (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Human, Rat**Predicted MW.:** 88 kDa**Subcellular Location:** Cytoplasm ,Nucleus**— VALIDATION IMAGES —**

Paraformaldehyde-fixed, paraffin embedded Human Pancreas; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with phospho-STAT3 (Tyr705) Monoclonal Antibody, Unconjugated(bsm-33301M) at 1:200 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with phospho-STAT3 (Tyr705) Monoclonal Antibody, Unconjugated(bsm-33301M) at 1:200 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.

— SELECTED CITATIONS —

- **[IF=8.109]** Ling Wu. et al. Calcium Channel Blocker Nifedipine Suppresses Colorectal Cancer Progression and Immune

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

- Escape by Preventing NFAT2 Nuclear Translocation. Cell Rep. 2020 Oct;33:108327 IF,IHC ;Human. 33113363
- **[IF=6.3]** Yijia Zhang. et al. β -Catenin mediated TAM phenotype promotes pancreatic cancer metastasis via the OSM/STAT3/LOXL2 axis. NEOPLASIA. 2025 Feb;60:101096 WB ;Human. 39740539
 - **[IF=4.414]** Wei Xiaojin. et al. BMP7 Attenuates Neuroinflammation after Spinal Cord Injury by Suppressing the Microglia Activation and Inducing Microglial Polarization Via the STAT3 Pathway. NEUROCHEM RES. 2023 Apr;;1-14 IF,WB ;Human. 37071344
 - **[IF=3.9]** Zhao JinFeng. et al. Synthesis, biological activities and mechanistic studies of C20-ketone pachysandra alkaloids as anti-hepatocellular carcinoma agents. MOL DIVERS. 2024 Aug;;1-21 ICC,WB ;Human. 39158620