

bsm-33227M**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

TBP Mouse mAb**— DATASHEET —****Host:** Mouse**Clonality:** Monoclonal**GeneID:** 6908**Target:** TBP**Purification:** affinity purified by Protein G**Concentration:** 1mg/ml

Storage: Size : 50ul/100ul
0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Size : 200ug (PBS only)
0.01M PBS
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes TBP, the TATA-binding protein. A distinctive feature of TBP is a long string of glutamines in the N-terminus. This region of the protein modulates the DNA binding activity of the C terminus, and modulation of DNA binding affects the rate of transcription complex formation and initiation of transcription. The number of CAG repeats encoding the polyglutamine tract is usually 25-42, and expansion of the number of repeats to 45-66 increases the length of the polyglutamine string and is associated with spinocerebellar ataxia 17, a neurodegenerative disorder classified as a polyglutamine disease. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2016]


Isotype: IgG**CloneNo.:** 6G8**SWISS:** P20226

Applications: **WB** (1:500-1000)
IHC-P (1:100-500)
IHC-F (1:100-500)
IF (1:100-500)


Reactivity: Human, Rat
(predicted: Mouse)

Predicted MW.: 38 kDa

Subcellular Location: Nucleus

— VALIDATION IMAGES —

Sample: U87MG(Human) Cell Lysate at 30 ug
Primary: Anti- TBP (bsm-33227M) at 1/1000



Paraformaldehyde-fixed, paraffin embedded (rat testis tissue); Antigen retrieval by boiling in

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

dilution Secondary: IRDye800CW Goat Anti-Mouse IgG at 1/20000 dilution Predicted band size: 38 kD Observed band size: 40 kD

sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TATA binding protein) Monoclonal Antibody, Unconjugated (bsm-33227M) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.