bs-0294M-Bio

[Secondary Antibodies]

Mouse Anti-Goat IgG H&L, Biotin conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –––––		400-901-9800
Host: Mouse Clonality: Polyclonal Target: Mouse Anti-Goat IgG Purification: affinity purified by Pr Concentration: 2.0 mg/ml		Applications: WB (1:1000-10000) IHC-P (1:100-500) IHC-F (1:100-1000) IF (1:100-1000) Flow-Cyt (1:100-1000) ICC/IF (1:100-1000) ELISA (1:1000-10000)
Storage: 10 mM TBS (pH=7.4) with 1% BSA, 0.03% Proclin300 and 50% glycerol. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Reactivity: Goat
serum with normal le is important for our c molecules are produ adaptive immune res functions; to bind to recruit other cells an variability of the IgG	gG), is one of the most abundant proteins evels between 8-17 mg/mL in adult blood lefence against microorganisms and the ced by B lymphocytes as a part of our sponse. The IgG molecule has two separa the pathogen that elicited the response a d molecules to destroy the antigen. The pool is generated by somatic recombinat pecificities in an individual at a given time be 1011 variants.	I. IgG and to tion