bs-4373R

[Primary Antibody]

ZNF710 Rabbit pAb




www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500) IF (1:100-500)
GenelD: 374655	SWISS: Q8N1W2	ICC/IF (1:50)
Target: ZNF710		Reactivity: Human, Mouse
Immunogen: KLH conjugated synthetic peptide derived from human ZNF710: 401-500/664.		(predicted: Rat, Pig, Sheep, Cow, Dog, Horse)
Purification: affinity purified by	Protein A	
Concentration: 1mg/ml		Predicted MW.: ^{75 kDa}
Storage: Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: ^{Nucleus}
its related pathway to this gene include	r Protein 710) is a Protein Coding gene. Among rs are Gene Expression. GO annotations related e nucleic acid binding and transcription cofactor ant paralog of this gene is ZNF366.	

— VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse colon); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (ZNF710) Polyclonal Antibody, Unconjugated (bs-4373R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (ZNF710) polyclonal Antibody, Unconjugated (bs-4373R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

- SELECTED CITATIONS -

• [IF=2.1] Li G et al. Overexpression of antisense long non-coding RNA ZNF710-AS1-202 promotes cell proliferation and inhibits apoptosis of clear cell renal cell carcinoma via regulation of ZNF710 expression. Mol Med Rep. 2020 Jun;21(6):2502-2512. WB ;Human. 32236626

ugatedconjugated Goat Anti-Rabbit IgG antibody atllowed by37°C for 90 minutes, DAPI (blue, C02-04002) wassp-0023)used to stain the cell nuclei.