

**bs-4382R**

**[ Primary Antibody ]**

## ZNF775 Rabbit pAb

**Bioss**  
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


### — DATASHEET —

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 285971	<b>SWISS:</b> Q96BV0	
<b>Target:</b> ZNF775		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human ZNF775: 61-160/537.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	<b>Reactivity:</b> Human, Mouse, Rat (predicted: Rabbit, Pig, Cow, Dog, Horse)	
<b>Background:</b> ZNF775 belongs to the krueppel C2H2 type zinc finger protein family. It contains 11 C2H2 type zinc fingers. ZNF775 may be involved in transcriptional regulation.	<b>Predicted MW.:</b> 60 kDa	
	<b>Subcellular Location:</b> Nucleus	


### — VALIDATION IMAGES —



Sample: Cerebrum (Mouse) Lysate at 40 ug  
Jurkat(Human) Cell Lysate at 30 ug  
Primary: Anti- ZNF775 (bs-4382R) at 1/1000 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 60 kD  
Observed band size: 60 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF775) Polyclonal Antibody, Unconjugated (bs-4382R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF775) Polyclonal Antibody, Unconjugated (bs-4382R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.