

bs-20237R**[Primary Antibody]****AMPK beta 1/PRKAB1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


DATASHEET**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 5564**SWISS:** Q9Y478**Target:** AMPK beta 1/PRKAB1**Immunogen:** KLH conjugated synthetic peptide derived from human AMPK beta 1 : 2-100/270.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: The protein encoded by this gene is a regulatory subunit of the AMP-activated protein kinase (AMPK). AMPK is a heterotrimer consisting of an alpha catalytic subunit, and non-catalytic beta and gamma subunits. AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. This subunit may be a positive regulator of AMPK activity. The myristoylation and phosphorylation of this subunit have been shown to affect the enzyme activity and cellular localization of AMPK. This subunit may also serve as an adaptor molecule mediating the association of the AMPK complex. [provided by RefSeq, Jul 2008].

Applications: IHC-P (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (2ug/Test)**ICC/IF** (1:100-500)**Reactivity:** Human, Rat
(predicted: Chicken)**Predicted
MW.:** 30 kDa**Subcellular
Location:** Cytoplasm ,Nucleus**VALIDATION IMAGES**


Paraformaldehyde-fixed, paraffin embedded (human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (AMPK beta 1/PRKAB1) Polyclonal Antibody, Unconjugated (bs-20237R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human kidney carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (AMPK beta 1/PRKAB1) Polyclonal Antibody, Unconjugated (bs-20237R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



MCF-7 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (AMPK beta 1/PRKAB1) polyclonal Antibody, Unconjugated (bs-20237R) 1:50, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Blank control (black line) :MCF-7. Primary
Antibody (green line): Rabbit Anti-AMPK beta
1/17 antibody (bs-20237R) Dilution:2ug/Test;
Secondary Antibody (white blue line) : Goat
anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Isotype
control (orange line) : Normal Rabbit IgG
Protocol The cells were fixed with 4% PFA
(10min at room temperature)and then
permeabilized with 90% ice-cold methanol for
20 min at -20°C, The cells were then incubated in
5%BSA to block non-specific protein-protein
interactions for 30 min at room temperature
.Cells stained with Primary Antibody for 30 min
at room temperature. The secondary antibody
used for 40 min at room temperature.
Acquisition of 20,000 events was performed.