## bs-1158R

## [ Primary Antibody ]

# Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# AGEs Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: AGEs

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** This gene encodes the most abundant protein in human blood.

This protein functions in the regulation of blood plasma colloid osmotic pressure and acts as a carrier protein for a wide range of endogenous molecules including hormones, fatty acids, and metabolites, as well as exogenous drugs. Additionally, this protein exhibits an esterase-like activity with broad substrate specificity. The encoded preproprotein is proteolytically processed to generate the mature protein. A peptide derived from this protein, EPI-X4, is an endogenous inhibitor of the CXCR4 chemokine

receptor. [provided by RefSeq, Jul 2016]

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)


Flow-Cyt (1ug/Test)

Reactivity: Human, Mouse, Rat

Predicted MW.: 62 k

Subcellular Location: Secreted


#### VALIDATION IMAGES




25 ug total protein per lane of various lysates (see on figure) probed with AGEs polyclonal antibody, unconjugated (bs-1158R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



25 ug total protein per lane of various lysates (see on figure) probed with AGEs polyclonal antibody, unconjugated (bs-1158R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Tissue/cell: Rat kidney tissue; 4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-AGEs Polyclonal Antibody, Unconjugated(bs-1158R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control: MCF7. Primary Antibody (green line): Rabbit Anti-AGEs antibody (bs-1158R) Dilution:  $1\mu g/10^{\circ}6$  cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody: Goat anti-rabbit IgG-AF647 Dilution:  $1\mu g/test$ . Protocol The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature

.Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

### - SELECTED CITATIONS -

- [IF=15.7] Zhang Zhongyu. et al. Galectin-3-integrin α5β1 phase separation disrupted by advanced glycation endproducts impairs diabetic wound healing in rodents. NAT COMMUN. 2025 Aug;16(1):1-23 IHC; Rat. 40775187
- [IF=14.593] Wanyi Zhou. et al. Glucose and MMP-9 dual-responsive hydrogel with temperature sensitive self-adaptive shape and controlled drug release accelerates diabetic wound healing. Bioact Mater. 2022 Jan;: IF; Rat. 10.1016/j.bioactmat.2022.01.004
- [IF=12.8] Yang Sheng. et al. Advanced glycation end products promote meniscal calcification by activating the mTOR-ATF4 positive feedback loop. EXP MOL MED. 2024 Mar;:1-16 IF,WB;Mouse,Human. 38424194
- [IF=10] Jun Ren. et al.Multitasking Asynchronous Collaborative Nanosystem for Diabetic Wound Healing Based on Hypoglycemic, Antimicrobial, and Angiogenesis-Promoting Effects.advanced healthcare materials.2025 Feb;14(4):e2403282. IHC; Mouse. 3968678
- [IF=7.8] Tianyu Dai. et al. Synovial advanced glycosylation end products aggravate periprosthetic infection in diabetes by upregulating Staphylococcus aureus RNAIII. J ORTHOP TRANSL. 2025 Jul;53:161 IHC,IF; Human,Rat. 40656895