

bs-10380R**[Primary Antibody]****Phospho-Smad1 (Ser465) Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 4086	SWISS: Q15797	IHC-F (1:100-500)
Target: Phospho-Smad1 (Ser465)		IF (1:100-500)
Immunogen: KLH conjugated Synthesised phosphopeptide derived from human Smad1 around the phosphorylation site of Ser465: SV(p-S)-NH ₂ .		Reactivity: Human (predicted: Mouse, Rat, Rabbit, Pig, Cow, Chicken, Horse)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 51 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Nucleus
Background: The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signals of the bone morphogenetic proteins (BMPs), which are involved in a range of biological activities including cell growth, apoptosis, morphogenesis, development and immune responses. In response to BMP ligands, this protein can be phosphorylated and activated by the BMP receptor kinase. The phosphorylated form of this protein forms a complex with SMAD4, which is important for its function in the transcription regulation. This protein is a target for SMAD-specific E3 ubiquitin ligases, such as SMURF1 and SMURF2, and undergoes ubiquitination and proteasome-mediated degradation. Alternatively spliced transcript variants encoding the same protein have been observed. [provided by RefSeq].		

— VALIDATION IMAGES —

Sample: Lane 1: Human Hela cell (UV) Lysates
 Lane 2: Human A431 cell (UV) Lysates Lane 3:
 Human U251 cell (UV) Lysates Lane 4: Human
 293T cell (TPA) Lysates Primary: Anti-Phospho-
 Smad1 (bs-10380R) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at
 1/20000 dilution Predicted band size: 51kDa
 Observed band size: 63kDa



Paraformaldehyde-fixed, paraffin embedded
 (Human brain glioma); Antigen retrieval by
 boiling in sodium citrate buffer (pH6.0) for
 15min; Block endogenous peroxidase by 3%
 hydrogen peroxide for 20 minutes; Blocking
 buffer (normal goat serum) at 37°C for 30min;
 Antibody incubation with (Histone H3 (mono
 methyl K4)) Polyclonal Antibody, Unconjugated
 (bs-2814R) at 1:400 overnight at 4°C, followed by
 operating according to SP Kit(Rabbit) (sp-0023)
 instructions and DAB staining.