

bs-13645R**[Primary Antibody]****IQSEC2 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

DATASHEET

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>GeneID: 23096</p> <p>Target: IQSEC2</p> <p>Immunogen: KLH conjugated synthetic peptide derived from human IQSEC2: 851-950/1478.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: IQSEC2 is a 1,478 amino acid protein that belongs to the BRAG family and contains one IQ domain, one PH domain and a SEC7 domain. Localizing to the cytoplasm, IQSEC2 is expressed in brain, kidney and small intestine, with weaker levels of expression in placenta, pancreas, ovary, prostate and liver. IQSEC2 is a component of the postsynaptic density at excitatory synapses, and interacts with ARF family members as a guanine nucleotide exchange factor. Through the activation of ARF substrates, IQSEC2 may play a crucial role in cytoskeletal and synaptic organization. The gene encoding IQSEC2 maps to the human X chromosome. Defects to the IQSEC2 gene have been linked to mental retardation X-linked type 1 (MRX1), a condition characterized by decreased intellectual function. IQSEC2 exists as three isoforms due to alternative splicing events.</p>	<p>Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)</p> <p>Reactivity: Mouse (predicted: Human, Rat, Pig, Sheep, Cow, Dog)</p> <p>Predicted MW.: 162 kDa</p> <p>Subcellular Location: Cytoplasm</p>
--	--

VALIDATION IMAGES

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (IQSEC2) Polyclonal Antibody, Unconjugated (bs-13645R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.