

bsm-33243M**[Primary Antibody]**

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

Transferrin Mouse mAb, Serum Loading Control**— DATASHEET —****Host:** Mouse**Clonality:** Monoclonal**GeneID:** 7018**Target:** Transferrin**Purification:** affinity purified by Protein G**Concentration:** 1mg/ml


Storage: Size : 50ul/100ul/500ul
0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Size : 200ug (PBS only)
0.01M PBS
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of this protein is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. This protein may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum. [provided by RefSeq, Sep 2009].


Isotype: IgG**CloneNo.:** 12A7**SWISS:** P02787**Applications:** WB (1:5000-10000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (2ug/Test)**Reactivity:** Human, Mouse, Rat, Cow

Predicted
MW.: 77 kDa


Subcellular
Location: Secreted

— VALIDATION IMAGES —


25 ug total protein per lane of various lysates (see on figure) probed with Transferrin monoclonal antibody, unconjugated (bsm-33243M) at 1:5000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded (human skin cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Transferrin) Monoclonal Antibody, Unconjugated (bsm-33243M) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Transferrin) Monoclonal Antibody, Unconjugated (bsm-33243M) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Mouse)(sp-0024) instructions and DAB staining.



Blank control (black line) :HepG2. Primary
Antibody (green line): Mouse Anti-
Transferrin(Serum Loading Control) antibody

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

(bsm-33243M) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-mouse IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Mouse IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.