

bs-2160R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

COL4A3/Tumstatin Rabbit pAb**— DATASHEET —**

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-600)
Clonality: Polyclonal		IHC-F (1:100-600)
GeneID: 1285	SWISS: Q01955	IF (1:100-600)
Target: COL4A3/Tumstatin		Reactivity: Mouse (predicted: Human, Rat)
Immunogen: KLH conjugated synthetic peptide derived from human Tumstatin / Collagen alpha-3(IV) chain: 1571-1670/1670.		
Purification: affinity purified by Protein A		Predicted MW.: 27/159 kDa
Concentration: 1mg/ml		Subcellular Location: Secreted ,Extracellular matrix ,Cell membrane
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Type IV collagen, the major structural component of basement membranes, is a multimeric protein composed of 3 alpha subunits. These subunits are encoded by 6 different genes, alpha 1 through alpha 6, each of which can form a triple helix structure with 2 other subunits to form type IV collagen. This gene encodes alpha 3. In the Goodpasture syndrome, autoantibodies bind to the collagen molecules in the basement membranes of alveoli and glomeruli. The epitopes that elicit these autoantibodies are localized largely to the non-collagenous C-terminal domain of the protein. A specific kinase phosphorylates amino acids in this same C-terminal region and the expression of this kinase is upregulated during pathogenesis. This gene is also linked to an autosomal recessive form of Alport syndrome. The mutations contributing to this syndrome are also located within the exons that encode this C-terminal region. Like the other members of the type IV collagen gene family, this gene is organized in a head-to-head conformation with another type IV collagen gene so that each gene pair shares a common promoter. [provided by RefSeq, Jun 2010]		

— VALIDATION IMAGES —

Paraformaldehyde-fixed, paraffin embedded (mouse lung tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (COL4A3) Polyclonal Antibody, Unconjugated (bs-2160R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

- **[IF=2.56]** Yasuda, Jumpei, et al. "T3 peptide, a fragment of tumstatin, stimulates proliferation and migration of cardiac

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

fibroblasts through activation of Akt signaling pathway." Naunyn-Schmiedeberg's Archives of Pharmacology(2017): 1-10.
WB ;="Rat". 28785775