

bs-2502R**[Primary Antibody]****HOXA10 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —


Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1ug/test) Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Cow, Dog, Horse) Predicted MW.: 42 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 3206	SWISS: P31260	
Target: HOXA10		
Immunogen: KLH conjugated synthetic peptide derived from human HOXA10: 311-410/410.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosome 7 and encodes a DNA-binding transcription factor that may regulate gene expression, morphogenesis, and differentiation. More specifically, it may function in fertility, embryo viability, and regulation of hematopoietic lineage commitment. Alternatively spliced transcript variants have been described. Read-through transcription also exists between this gene and the downstream homeobox A9 (HOXA9) gene. [provided by RefSeq, Mar 2011].		

— VALIDATION IMAGES —


Paraformaldehyde-fixed, paraffin embedded (rat testis tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HOXA10) Polyclonal Antibody, Unconjugated (bs-2502R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse testis tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HOXA10) Polyclonal Antibody, Unconjugated (bs-2502R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control (black line) :NIH/3T3. Primary Antibody (green line): Rabbit Anti-HOXA10 antibody (bs-2502R) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



Blank control (black line) :A431. Primary Antibody (green line): Rabbit Anti-HOXA10 antibody (bs-2502R) Dilution:1ug/Test; Secondary Antibody (white blue line) : Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test. Isotype control (orange line) : Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=5.763]** You Shuang. et al. Recombinant humanized collagen remodels endometrial immune microenvironment of chronic endometritis through macrophage immunomodulation. REGEN BIOMATER. 2023 Apr;; IHC ;Rat. 37122820
- **[IF=4.501]** Fang Shi. et al. Phosphatidylinositol 3-Kinase/Protein Kinase B/Mammalian Target of the Rapamycin Pathway-Related Protein Expression in Lung Squamous Cell Carcinoma and Its Correlation with Lymph Node Metastasis.. J ONCOL. 2022 Aug;2022:4537256-4537256 WB ;Human. 36052284
- **[IF=5.131]** Yu Tang. et al. hUMSCs Restore Uterine Function by Inhibiting Endometrial Fibrosis via Regulation of the MMP-9/TIMP-1 Ratio in CDDP-Induced Injury Rats. STEM CELLS INT. 2023;2023:8014052 IHC ;Rat. 36994440
- **[IF=4.8]** Yang Qian. et al. Human Umbilical Cord Mesenchymal Stem Cells Promote Anti-Inflammation and Angiogenesis by Targeting Macrophages in a Rat Uterine Scar Model. STEM CELL REV. 2024 May;;1-14 IHC ;Rat. 38703310
- **[IF=3.8]** Rui Ren. et al. Developmental exposure to perfluorooctane sulfonate(PFOS) impairs the endometrial receptivity.SCIENTIFIC REPORTS.2025 Jan 11;15(1):1747. IF ;Rabbit. 39799182