bs-5521R

[Primary Antibody]

BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

IHC-F (1:100-500)

Mouse, Rabbit, Pig, Dog)

IF (1:100-500)

Reactivity: Rat (predicted: Human,

147/319 kDa

Applications: IHC-P (1:100-500)

Predicted

MW.:

Subcellular Nucleus

phospho-NF1 (Ser2817) Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 4763 **SWISS:** P21359

Target: NF1 (Ser2817)

Immunogen: KLH conjugated Synthesised phosphopeptide derived from human

NF1 around the phosphorylation site of Ser2817: HG(p-S)AS.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.


Background: Neurofibromin is a product of the tumor suppressor gene,

Neurofibromatosis type I. Neurofibromin is known to have GTPase activity that modulates the ras pathway. The absence of or

alteration of the neurofibromin protein may lead to

Neurofibromatosis disease. This protein has not been purified, therefore, most of the information regarding this protein has been deduced from homology analysis of its gene sequence.

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat testis tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NF1(Ser2817)) Polyclonal Antibody, Unconjugated (bs-5521R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NF1(Ser2817)) Polyclonal Antibody, Unconjugated (bs-5521R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.