## bs-6088R

## [ Primary Antibody ]

# Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# RASAL1 Rabbit pAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**GenelD:** 8437 **SWISS:** 095294

Target: RASAL1

**Immunogen:** KLH conjugated synthetic peptide derived from human RASAL1:

451-550/805.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Ras GTPase activating-like protein (RASAL) or RASAL1 is a member

of the GAP1 family, and a Ca2+ sensor responding in-phase to repetitive Ca2+ signals by associating with the plasma membrane and deactivating Ras. It contains a conserved domain structure comprising N-terminal tandem C2 domains, a highly conserved central RasGAP domain, and a C-terminal pleckstrin-homology domain that is associated with a Bruton's tyrosine kinase motif. RASAL, like Ca2+-promoted Ras inactivator (CAPRI, or RASAL4), is a cytosolic protein that undergoes a rapid translocation to the plasma membrane in response to receptor-mediated elevation in the concentration of intracellular free Ca2+, a translocation that activates its ability to function as a RasGAP. However, unlike RASAL4, RASAL undergoes an oscillatory translocation to the plasma membrane that occurs in synchrony with repetitive Ca2+

spikes.

Applications: IHC-P (1:400-800)

**IHC-F** (1:400-800) **IF** (1:100-500)

Reactivity: Human, Rat


(predicted: Mouse, Sheep,

Cow, Dog)


Predicted MW.: 90 kDa

Subcellular Cytoplasm

#### VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (human gastric carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RASAL1) Polyclonal Antibody, Unconjugated (bs-6088R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Tissue/cell: rat brain tissue;4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min;
Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-RASAL1
Polyclonal Antibody, Unconjugated(bs-6088R) 1:200, overnight at 4°C; The secondary antibody was Goat Anti-Rabbit IgG, Cy3
conjugated(bs-0295G-Cy3)used at 1:200 dilution for 40 minutes at 37°C. DAPI(5ug/ml,blue,C-0033) was used to stain the cell nuclei

### — SELECTED CITATIONS ———

• [IF=3.06] Hong, Yan, et al. "Paridis Rhizoma Sapoinins attenuates liver fibrosis in rats by regulating the expression of RASAL1/ERK1/2 signal pathway." Journal of Ethnopharmacology (2016). IHC; Rat. 27396351