


bsm-60157M**[Primary Antibody]****Mono-Methyl-Histone H4 (Arg3) Mouse mAb****BioSS**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —**Host:** Mouse**Isotype:** IgG**Clonality:** Monoclonal**CloneNo.:** D2G9**Target:** Mono-Methyl-Histone H4 (Arg3)**Purification:** Antigen affinity purification**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. [provided by RefSeq, Jul 2008]**Applications:** WB (1:500-1:2000)**Reactivity:** Human (predicted: Mouse, Rat, Monkey)**Predicted MW.:** 11 kDa**Subcellular Location:** Nucleus**— VALIDATION IMAGES —**

Blocking buffer: 5% NFDM/TBST Primary ab
dilution: 1:2000 Primary ab incubation
condition: 2 hours at room temperature
Secondary ab: Goat Anti-Mouse IgG H&L (HRP)
Lysate: HeLa, C2C12, COS-7, BRL, N2a Protein
loading quantity: 20 µg Exposure time: 60 s
Predicted MW: 11 kDa Observed MW: 11 kDa