

**bsm-60142M****[ Primary Antibody ]****BioSS**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**Asymmetric Di-Methyl-Histone H4 (Arg3) Mouse mAb****— DATASHEET —**

<b>Host:</b> Mouse	<b>Isotype:</b> IgG	<b>Applications:</b> WB (1:500-1:2000)  <b>Reactivity:</b> Human (predicted: Mouse, Rat, Monkey)  <b>Subcellular Location:</b> Nucleus
<b>Clonality:</b> Monoclonal	<b>CloneNo.:</b> D5C7	
<b>Target:</b> Asymmetric Di-Methyl-Histone H4 (Arg3)		
<b>Purification:</b> Antigen affinity purification		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. [provided by RefSeq, Jul 2008]		

**— VALIDATION IMAGES —**

Blocking buffer: 5% NFDm/TBST Primary ab  
dilution: 1:2000 Primary ab incubation  
condition: 2 hours at room temperature  
Secondary ab: Goat Anti-Mouse IgG H&L (HRP)  
Lysate: HeLa, C2C12, COS-7 Protein loading  
quantity: 20 µg Exposure time: 30 s Predicted  
MW: 11 kDa Observed MW: 11 kDa