


bs-42280R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

PPARGC1A Rabbit pAb**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 10891**SWISS:** Q9UBK2**Target:** PPARGC1A**Immunogen:** Recombinant human PPARGC1A: 11-280/798.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.**Background:** The protein encoded by this gene is a transcriptional coactivator that regulates the genes involved in energy metabolism. This protein interacts with PPARGgamma, which permits the interaction of this protein with multiple transcription factors. This protein can interact with, and regulate the activities of cAMP response element binding protein (CREB) and nuclear respiratory factors (NRFs). It provides a direct link between external physiological stimuli and the regulation of mitochondrial biogenesis, and is a major factor that regulates muscle fiber type determination. This protein may be also involved in controlling blood pressure, regulating cellular cholesterol homoeostasis, and the development of obesity (referenced from entrez gene).**Applications:** **WB** (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**ICC/IF** (1:50-100)**Reactivity:** Human (predicted: Mouse, Rat, Goat)**Predicted MW.:** 88 kDa**Subcellular Location:** Nucleus**— VALIDATION IMAGES —**


Sample: Lane 1: Recombinant human PPARGC1A, N-Trx-His(bs-42280P) Primary: Anti-PPARGC1A (bs-42280R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 88 kDa
Observed band size: 61 kDa



Paraformaldehyde-fixed, paraffin embedded (human kidney carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (PPARGC1A) Polyclonal Antibody, Unconjugated (bs-42280R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (PPARGC1A) Polyclonal Antibody, Unconjugated (bs-42280R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



A431 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

min; Antibody incubation with (PPARGC1A)
polyclonal Antibody, Unconjugated (bs-42280R)
1:50, 90 minutes at 37°C; followed by a
conjugated Goat Anti-Rabbit IgG antibody at
37°C for 90 minutes, DAPI (blue, C02-04002) was
used to stain the cell nuclei.