## bs-33011R

## [ Primary Antibody ]

## BIOSS

RFP Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: RFP

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Fluorescent proteins have become a useful and ubiquitous tool for

making chimeric proteins, where they function as a fluorescent protein tag. Typically they tolerate N- and C-terminal fusion to a broad variety of proteins. They have been expressed in most known cell types and are used as a noninvasive fluorescent marker in living cells and organisms. They enable a wide range of applications where they have functioned as a cell lineage tracer, reporter of gene expression, or as a measure of protein-protein interactions. Red Fluorescent Protein (RFP) is a versatile biological marker for monitoring physiological processes, visualizing protein localization, and detecting transgenic expression in vivo. RFP can be excited by the 488 nm or 532 nm laser line and is optimally

detected at 588 nm.


Applications: WB (1:500-2000)

IHC-P (1:200-1000)
IHC-F (1:200-1000)
IF (1:200-1000)
Flow-Cyt (1ug/Test)
ICC/IF (1:100-500)
ELISA (1:5000-10000)

Reactivity: RFP Tag

Subcellular Cytoplasm ,Nucleus

## VALIDATION IMAGES -



Sample: Lane 1: Recombinant RFP protein, His Lane 2: Tags Positive Control Whole Cell Lysate, C-His Primary: Anti-RFP (bs-33011R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: kDa Observed band size: 30,44 kDa