bs-0377R-APC

[Secondary Antibodies]

Rabbit Anti-Mouse IgG Fc, APC conjugated



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –––––		400-901-9800
Host: Rabbit Clonality: Polyclonal	lsotype: IgG	Applications: IF (1:100-1000) Flow-Cyt (1:100-1000)
Target: Rabbit Anti-Mouse IgG Fc Purification: affinity purified by Protein A		ICC/IF (1:100-1000) Excitation Spectrum: 650nm Emission spectrum: 660nm Reactivity: Mouse
glycerol.	vith 1% BSA, 0.02% Proclin300 and 5 year. Avoid repeated freeze/thaw cy	
serum with normal le is important for our d molecules are produc adaptive immune res functions; to bind to recruit other cells and variability of the IgG p	gG), is one of the most abundant pro- vels between 8-17 mg/mL in adult bl efence against microorganisms and ed by B lymphocytes as a part of our ponse. The IgG molecule has two sep the pathogen that elicited the respor d molecules to destroy the antigen. T pool is generated by somatic recomb ecificities in an individual at a given be 1011 variants.	ood. IgG the r parate nse and to he ination