

**bs-4404R****[ Primary Antibody ]****NUP188 Rabbit pAb**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**— DATASHEET —**

<b>Host:</b> Rabbit	<b>Isotype:</b> IgG	<b>Applications:</b> <b>IHC-P</b> (1:100-500) <b>IHC-F</b> (1:100-500) <b>IF</b> (1:100-500) <b>ICC/IF</b> (1:100-500)  <b>Reactivity:</b> (predicted: Human, Mouse, Rat, Pig, Sheep, Cow, Chicken, Dog, Horse)  <b>Predicted MW.:</b> 196 kDa  <b>Subcellular Location:</b> Nucleus
<b>Clonality:</b> Polyclonal		
<b>GeneID:</b> 23511	<b>SWISS:</b> Q5SRE5	
<b>Target:</b> NUP188		
<b>Immunogen:</b> KLH conjugated synthetic peptide derived from human NUP188: 671-770/1749.		
<b>Purification:</b> affinity purified by Protein A		
<b>Concentration:</b> 1mg/ml		
<b>Storage:</b> Preservative: 0.02% Proclin300, Constituents: 1% BSA, 0.01M PBS, pH7.4. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
<b>Background:</b> The nuclear pore complex (NPC) is found on the nuclear envelope and forms a gateway that regulates the flow of proteins and RNAs between the cytoplasm and nucleoplasm. The NPC is comprised of approximately 30 distinct proteins collectively known as nucleoporins. Nucleoporins are pore-complex-specific glycoproteins which often have cytoplasmically oriented O-linked N-acetylglucosamine residues and numerous repeats of the pentapeptide sequence XFXFG. However, the nucleoporin protein encoded by this gene does not contain the typical FG repeat sequences found in most vertebrate nucleoporins. This nucleoporin is thought to form part of the scaffold for the central channel of the nuclear pore. [provided by RefSeq, Jan 2013]		