

PKC zeta Rabbit pAb

Catalog Number: bs-10399R

Target Protein: PKC zeta

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ICC/IF (1:100-500), ELISA (1:5000-10000)

Reactivity: (predicted:Human, Mouse, Rat, Rabbit, Pig, Sheep, Cow, Dog, Horse)

Predicted MW: 66 kDa

Entrez Gene: 5590

Swiss Prot: Q05513

Source: KLH conjugated synthetic peptide derived from human PKC zeta: 61-160/592.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Protein kinase C (PKC) zeta is a member of the PKC family of serine/threonine kinases which are involved in a variety of cellular processes such as proliferation, differentiation and secretion. Unlike the classical PKC isoenzymes which are calcium-dependent, PKC zeta exhibits a kinase activity which is independent of calcium and diacylglycerol but not of phosphatidylserine. Furthermore, it is insensitive to typical PKC inhibitors and cannot be activated by phorbol ester. Unlike the classical PKC isoenzymes, it has only a single zinc finger module. These structural and biochemical properties indicate that the zeta subspecies is related to, but distinct from other isoenzymes of PKC. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008].

PRODUCT SPECIFIC PUBLICATIONS

[IF=3.365] Xin Li. et al. Hengshun Aromatic Vinegar Ameliorates Vascular Endothelial Injury via Regulating PKC ζ -Mediated Oxidative Stress and Apoptosis. Front Nutr. 2021; 8: 635232 IF ; Rat . 34124116