bsm-60878R

[Primary Antibody]

www.bioss.com.cn

sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

OLIG1 Recombinant Rabbit mAb

DATASHEET -

Host: Rabbit Isotype: IgG Clonality: Recombinant CloneNo.: 2B3 **GenelD: 116448** SWISS: Q8TAK6

Target: OLIG1

Immunogen: A synthesized peptide derived from human Oligo1: 200-271/271.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS(pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Shipped at 4°C. Store at -20 °C for one year. Avoid repeated

freeze/thaw cycles.

Background: The oligodendrocyte lineage-specific basic helix-loop-helix (OLIG) family of transcription factors include OLIG1-OLIG3, which differ in tissue expression. OLIG1 and OLIG2 are specifically expressed in nervous tissue as gene regulators of oligodendrogenesis. OLIG2 is more widely expressed in embryonic brain than OLIG1, while OLIG3 is primarily expressed in non-neural tissues. OLIG1 and OLIG2 interact with the Nkx-2.2 homeodomain protein, which is responsible for directing ventral neuronal patterning in response to graded Sonic hedgehog signaling in the embryonic neural tube. These interactions between OLIG proteins and Nkx-2.2 appear to promote the formation of alternate cell types by inhibiting V3 interneuron development. OLIG1 and OLIG2 are abundantly expressed in oligodendroglioma and nearly absent in astrocytomas. Therefore, OLIG proteins are candidates for molecular markers of human glial brain tumors, which are the most common primary malignancies of the human brain.

Applications: WB (1:500-2000)


Reactivity: Human (predicted: Mouse,

Rat)

Predicted 28 kDa MW.:

Subcellular Nucleus

VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with OLIG1 monoclonal antibody, unconjugated (bsm-60878R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.