

bs-6331R**[Primary Antibody]****NET1 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 10276**SWISS:** Q7Z628**Target:** NET1**Immunogen:** KLH conjugated synthetic peptide derived from human NET1: 501-596/596.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene is part of the family of Rho guanine nucleotide exchange factors. Members of this family activate Rho proteins by catalyzing the exchange of GDP for GTP. The protein encoded by this gene interacts with RhoA within the cell nucleus and may play a role in repairing DNA damage after ionizing radiation. Pseudogenes of this gene are located on the long arms of chromosomes 1, 7 and 18. Alternative splicing results in multiple transcript variants that encode different protein isoforms.**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (2ug/Test)**ICC/IF** (1:25)**Reactivity:** Human, Mouse
(predicted: Rat, Cow,
Chicken, Dog, Horse)**Predicted
MW.:** 65 kDa**Subcellular
Location:** Cytoplasm ,Nucleus**— VALIDATION IMAGES —**


Sample: Jurkat(Human) Cell Lysate at 30 ug
 Primary: Anti-NET1 (bs-6331R) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 65 kD
 Observed band size: 65 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NET1) Polyclonal Antibody, Unconjugated (bs-6331R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



HepG2 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (NET1) polyclonal Antibody, Unconjugated (bs-6331R) 1:25, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control (black line) :HepG2. Primary
 Antibody (green line): Rabbit Anti-NET1 antibody
 (bs-6331R) Dilution:2ug/Test; Secondary
 Antibody (white blue line) : Goat anti-rabbit
 IgG-FITC Dilution: 0.5ug/Test. Isotype control
 (orange line) : Normal Rabbit IgG Protocol The

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.