

**bs-0766R****[ Primary Antibody ]****CD4 Rabbit pAb****Bioss**  
**ANTIBODIES**

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 12504**SWISS:** P06332**Target:** CD4**Immunogen:** KLH conjugated synthetic peptide derived from the middle of mouse CD4: 231-330/457. < Extracellular >**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

**Background:** This gene encodes a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex class II antigens and is also a receptor for the human immunodeficiency virus. This gene is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified in this gene. [provided by RefSeq, Aug 2010].**Applications:** WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Reactivity:** Human, Mouse, Rat**Predicted MW.:** 48 kDa**Subcellular Location:** Cell membrane**— VALIDATION IMAGES —**


Sample: Lane 1: Human HL-60 cell lysates Lane 2: Human MOLT4 cell lysates Lane 3: Human Jurkat cell lysates Lane 4: Human Raji cell lysates Primary: Anti-CD4 (bs-0766R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 48 kDa> Observed band size: 60 kDa




Sample: U937(Human) Cell Lysate at 30 ug Primary: Anti- CD4 (bs-0766R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 48 kD Observed band size: 55 kD



Paraformaldehyde-fixed, paraffin embedded (rat spleen); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CD4) Polyclonal Antibody, Unconjugated (bs-0766R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Tissue/cell: mouse lymphoma tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (



Tissue/cell: rat lung tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-CD4 Polyclonal Antibody, Unconjugated(bs-0766R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

0.01M, pH 6.0 ), Boiling bathing for 15min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-CD4(mouse, rat) Polyclonal Antibody, Unconjugated(bs-0766R) 1:200, overnight at 4°C; The secondary antibody was Goat Anti-Rabbit IgG, Cy3 conjugated(bs-0295G-Cy3)used at 1:200 dilution for 40 minutes at 37°C. DAPI(5ug/ml,blue,C-0033) was used to stain the cell nuclei

## — SELECTED CITATIONS —

- **[IF=15.1]** Lei Wang. et al. STING Agonist-Loaded Nanoparticles Promotes Positive Regulation of Type I Interferon-Dependent Radioimmunotherapy in Rectal Cancer. ADV SCI. 2023 Dec;;2307858 IF ;Mouse. 38063844
- **[IF=15.304]** Jinbo Li. et al. Autophagy inhibition recovers deficient ICD-based cancer immunotherapy. BIOMATERIALS. 2022 Aug;287:121651 IF ;Mouse. 35777331
- **[IF=8.806]** Li TF et al. Dendritic cell-mediated delivery of doxorubicin-polyglycerol-nanodiamond composites elicits enhanced anti-cancer immune response in glioblastoma.Biomaterials. 2018 Oct;181:35-52. IHC ;Human. 30071380
- **[IF=7.727]** Jinbo Li. et al. Low dose shikonin and anthracyclines coloaded liposomes induce robust immunogenetic cell death for synergistic chemo-immunotherapy. J Control Release. 2021 Jul;335:306 IHC ;Mouse. 34081995
- **[IF=8.39]** Wang, Raymond M., et al. "Humanized mouse model for assessing the human immune response to xenogeneic and allogeneic decellularized biomaterials." Biomaterials (2017). IHC ;="Mouse". 28334641