### bs-15398R

## [ Primary Antibody ]

# HHV11 ICP22 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Target: HHV11 ICP22

**Immunogen:** KLH conjugated synthetic peptide derived from human HHV-1:

101-200/420.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Herpes simplex type 1 (HSV-1) belongs to a family that includes

HSV-2, Epstein-Barr virus (EBV) and Varicella zoster (chicken pox) virus amongst others. HSV-1 and HSV-2 are extremely difficult to distinguish from each other. Members of this family have a

characteristic virion structure. The double stranded DNA genome is contained within an icosahedral capsid embedded in a proteinaceous layer (tegument) and surrounded by a lipid envelope, derived from the nuclear membrane of the last host, which is decorated with virus-specific glycoproteins spikes. These viruses are capable of entering a latent phase where the host

shows no visible sign of infection and levels of infectious agent become very low. During the latent phase the viral DNA is integrated into the genome of the host cell.

Applications: WB (1:500-2000)

**IHC-P** (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) ICC/IF (1:100-500) **ELISA** (1:5000-10000)

Reactivity: (predicted: Herpes Simplex

Virus)

Predicted MW.: 46 kDa

Subcellular Location: Cytoplasm ,Nucleus

### - SELECTED CITATIONS -

• [IF=7.5] Hansong Qi. et al. ICP22-defined condensates mediate RNAPII deubiquitylation by UL36 and promote HSV-1 transcription. CELL REP. 2024 Oct;43: WB; Human. 39383039