

bs-7106R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

NUPR1 Rabbit pAb**DATASHEET**

Host: Rabbit Clonality: Polyclonal GeneID: 26471 Target: NUPR1 Immunogen: KLH conjugated synthetic peptide derived from human NUPR1/p8: 31-82/82. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Could participate in the response to proapoptotic stimuli and promotes cellular growth in a way that helps the tissue counteract diverse injuries. May contribute to the metastatic phenotype. Tissue specificity: Highly expressed in pancreas.	Isotype: IgG SWISS: O60356	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ELISA (1:5000-10000) Reactivity: Human, Mouse (predicted: Rat, Sheep, Cow, Dog, Horse) Predicted MW.: 9 kDa Subcellular Location: Cytoplasm ,Nucleus
--	---	--

VALIDATION IMAGES

Sample: Lane 1: Human L-O2 cell lysates Lane 2: Human U87MG cell lysates Primary: Anti-NUPR1 (bs-7106R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 9 kDa Observed band size: 11 kDa



Paraformaldehyde-fixed, paraffin embedded (Mouse pancreas); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (NUPR1) Polyclonal Antibody, Unconjugated (bs-7106R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

SELECTED CITATIONS

- **[IF=24.633]** Yanxin Li. et al. Decoding the temporal and regional specification of microglia in the developing human brain. Cell Stem Cell. 2022 Mar; IF ;Mouse,Human. 35245443
- **[IF=7.126]** Narzt et al. A novel role for NUPR1 in the keratinocyte stress response to UV oxidized phospholipids. (2018) Redox.Biol. 20:467-482 IF ;Human. 30466060
- **[IF=6.1]** Zhou Jie. et al. Nupr1-mediated vascular smooth muscle cell phenotype transformation involved in methamphetamine induces pulmonary hypertension. CELL BIOL TOXICOL. 2024 Dec;40(1):1-20 IHC,IF ;Mouse. 38347241
- **[IF=6.1]** Liao Taiyang. et al. Suppression of NUPR1 in fibroblast-like synoviocytes reduces synovial fibrosis via the Smad3 pathway. J TRANSL MED. 2024 Dec;22(1):1-14 IF,WB ;Mouse. 39090667
- **[IF=5.391]** Shiraki M et al. Deficiency of stress-

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

associated gene Nupr1 increases bone volume by attenuating differentiation of osteoclasts and enhancing differentiation of osteoblasts. FASEB J. 2019 May 8:fj201802322RR. IF ;MOUSE. 31067083