

bs-1202R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

TRADD Rabbit pAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		IHC-P (1:100-500)
GeneID: 71609	SWISS: Q3U0V2	IHC-F (1:100-500)
Target: TRADD		IF (1:100-500)
Immunogen: KLH conjugated synthetic peptide derived from the middle of mouse TRADD: 111-230/310.		Reactivity: Human, Mouse, Rat (predicted: Rabbit)
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Predicted MW.: 34 kDa
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Subcellular Location: Cytoplasm ,Nucleus
Background: TNFR1 associated DEATH domain protein; TNFR1-associated DEATH domain protein; TNFRSF1A associated via death domain; TNFRSF1A-associated via death domain; tradd; TRADD_HUMAN; Tumor necrosis factor receptor type 1 associated DEATH domain protein; Tumor necrosis factor receptor type 1-associated DEATH domain protein.		

— VALIDATION IMAGES —


Sample: Lane1: Colon carcinoma (Human) Lysate at 30 ug Lane2: Brain (Mouse) Lysate at 30 ug
Primary: Anti-TRADD (bs-1202R) at 1:200 dilution; Secondary: HRP conjugated Goat Anti-Rabbit IgG (bs-0295G-HRP) at 1:3000 dilution;
Predicted band size: 35kD Observed band size: 34kD



Sample: Cerebrum (Mouse) Lysate at 40 ug
Primary: Anti-TRADD (bs-1202R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 34 kD



Tissue/cell: rat pancreas tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-TRADD Polyclonal Antibody, Unconjugated (bs-1202R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody (SP-0023) and DAB (C-0010) staining



Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-TRADD Polyclonal Antibody,

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Unconjugated(bs-1202R) 1:200, overnight at 4°C,
followed by conjugation to the secondary
antibody(SP-0023) and DAB(C-0010) staining

— SELECTED CITATIONS —

- **[IF=2.37]** Chen, Zhiqiang, et al. "Hydrogen sulfide protects against TNF- α induced neuronal cell apoptosis through miR-485-5p/TRADD signaling." Biochemical and Biophysical Research Communications (2016). WB ;="Human". 27562714