

TRADD Rabbit pAb

Catalog Number: bs-1202R

Target Protein: TRADD

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Rabbit)

Predicted MW: 34 kDa

Entrez Gene: 71609

Swiss Prot: Q3U0V2

Source: KLH conjugated synthetic peptide derived from the middle of mouse TRADD: 111-230/310.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: TNFR1 associated DEATH domain protein; TNFR1-associated DEATH domain protein; TNFRSF1A associated via death domain; TNFRSF1A-associated via death domain; tradd; TRADD_HUMAN; Tumor necrosis factor receptor type 1 associated DEATH domain protein; Tumor necrosis factor receptor type 1-associated DEATH domain protein.


VALIDATION IMAGES




Sample: Lane1: Colon carcinoma (Human) Lysate at 30 ug Lane2: Brain (Mouse) Lysate at 30 ug Primary: Anti-TRADD (bs-1202R) at 1:200 dilution; Secondary: HRP conjugated Goat Anti-Rabbit IgG (bs-0295G-HRP) at 1:3000 dilution; Predicted band size : 35kD Observed band size : 34kD




Sample: Jurkat(Human) Cell Lysate at 30 ug MCF-7(Human) Cell Lysate at 30 ug Primary: Anti- TRADD (bs-1202R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 34 kD



Sample: Cerebrum (Mouse) Lysate at 40 ug Primary: Anti- TRADD (bs-1202R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 34 kD



Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-TRADD Polyclonal Antibody, Unconjugated(bs-1202R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat pancreas tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-TRADD Polyclonal Antibody, Unconjugated(bs-1202R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

PRODUCT SPECIFIC PUBLICATIONS

[IF=2.37] Chen, Zhiqiang, et al. "Hydrogen sulfide protects against TNF- α induced neuronal cell apoptosis through miR-485-5p/TRADD signaling." Biochemical and Biophysical Research Communications (2016). WB ; ="Human" . 27562714