### bs-1202R

#### [ Primary Antibody ]

## www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# TRADD Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 71609 **SWISS:** Q3U0V2

Target: TRADD

Immunogen: KLH conjugated synthetic peptide derived from the middle of

mouse TRADD: 111-230/310.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: TNFR1 associated DEATH domain protein; TNFR1-associated

DEATH domain protein; TNFRSF1A associated via death domain; TNFRSF1A-associated via death domain; tradd; TRADD\_HUMAN; Tumor necrosis factor receptor type 1 associated DEATH domain protein; Tumor necrosis factor receptor type 1-associated DEATH

domain protein.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Human, Mouse, Rat


(predicted: Rabbit)

Predicted

34 kDa

**Subcellular Location:** Cytoplasm ,Nucleus


#### VALIDATION IMAGES




Sample: Lane1: Colon carcinoma (Human)Lysate at 30 ug Lane2:Brain(Mouse) Lysate at 30 ug Primary: Anti-TRADD (bs-1202R) at 1:200 dilution; Secondary: HRP conjugated Goat Anti-Rabbit IgG(bs-0295G-HRP) at 1: 3000 dilution; Predicted band size: 35kD Observed band size: 34kD



Sample: Cerebrum (Mouse) Lysate at 40 ug Primary: Anti- TRADD (bs-1202R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 34 kD Observed band size: 34 kD



Tissue/cell: rat pancreas tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-TRADD Polyclonal Antibody, Unconjugated(bs-1202R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat kidney tissue; 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-TRADD Polyclonal Antibody,

Unconjugated(bs-1202R) 1:200, overnight at 4°C
followed by conjugation to the secondary
antibody(SP-0023) and DAB(C-0010) staining

							т			

• [IF=2.37] Chen, Zhiqiang, et al. "Hydrogen sulfide protects against TNF- $\alpha$  induced neuronal cell apoptosis through miR-485-5p/TRADD signaling." Biochemical and Biophysical Research Communications (2016). WB; Human. 27562714