

bs-0152R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

TTR/TTR/Prealbumin Rabbit pAb**DATASHEET**

Host: Rabbit Clonality: Polyclonal GeneID: 7276 Target: TTR/Prealbumin Immunogen: KLH conjugated synthetic peptide derived from human Transthyretin: 51-147/147. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: This gene encodes transthyretin, one of the three prealbumins including alpha-1-antitrypsin, transthyretin and orosomucoid. Transthyretin is a carrier protein; it transports thyroid hormones in the plasma and cerebrospinal fluid, and also transports retinol (vitamin A) in the plasma. The protein consists of a tetramer of identical subunits. More than 80 different mutations in this gene have been reported; most mutations are related to amyloid deposition, affecting predominantly peripheral nerve and/or the heart, and a small portion of the gene mutations is non-amyloidogenic. The diseases caused by mutations include amyloidotic polyneuropathy, euthyroid hyperthyroxinaemia, amyloidotic vitreous opacities, cardiomyopathy, oculoleptomeningeal amyloidosis, meningocerebrovascular amyloidosis, carpal tunnel syndrome, etc. [provided by RefSeq]	Isotype: IgG Applications: WB (1:500-2000) Reactivity: Human, Mouse, Rat Predicted MW.: 14 kDa Subcellular Location: Secreted ,Cytoplasm
--	---

VALIDATION IMAGES

Sample: Lane 1: Mouse Serum Lane 2: Rat Serum
Lane 3: Human Plasma Lane 4: Human Serum
Primary: Anti-TTR/Prealbumin (bs-0152R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 14 kDa Observed band size: 14 kDa

SELECTED CITATIONS

- **[IF=4.5]** Yan Shu. et al. Bioaccumulation and Thyroid Endocrine Disruption of 2-Ethylhexyl Diphenyl Phosphate at Environmental Concentration in Zebrafish Larvae. AQUAT TOXICOL. 2023 Dec;;106815 WB ;Zebrafish. 10.1016/j.aquatox.2023.106815
- **[IF=4.872]** Dong X et al. PM2.5 disrupts thyroid hormone homeostasis through activation of the hypothalamic-pituitary-thyroid (HPT) axis and induction of hepatic transthyretin in female rats 2.5Ecotoxicol Environ Saf.2021 Jan 15;208:111720. IHC,WB ;Rat. 33396051
- **[IF=4.1]** Herrick-Davis, Katharine, et al. "Native Serotonin 5-HT2C Receptors are Expressed as Homodimers on the

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

- Apical Surface of Choroid Plexus Epithelial Cells." Molecular Pharmacology (2015): mol-114. Other ;="Rat". 25609374
- **[IF=4.223]** Dong, Xinwen. et al. Protective effects of curcumin against thyroid hormone imbalance after gas explosion-induced traumatic brain injury via activation of the hypothalamic-pituitary-thyroid axis in male rats. ENVIRON SCI POLLUT R. 2022 May;;1-13 WB,IHC ;Rat. 35641736
 - **[IF=3.738]** Zhou, Can-Can. et al. Lead Exposure in Developmental Ages Promotes A β Accumulation by Disturbing A β Transportation in Blood-Cerebrospinal Fluid Barrier/Blood–Brain Barriers and Impairing A β Clearance in the Liver. 2021 Nov 17 IHC ;Rat. 34787833