

bs-7111R**[Primary Antibody]****EDA2R Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500)
Clonality: Polyclonal		IHC-F (1:100-500)
GeneID: 60401	SWISS: Q9HAV5	IF (1:100-500)
Target: EDA2R		Reactivity: Human, Mouse (predicted: Rat)
Immunogen: KLH conjugated synthetic peptide derived from human EDA2R/TNFRSF27: 51-150/297.		
Purification: affinity purified by Protein A		Predicted MW.: 33 kDa
Concentration: 1mg/ml		Subcellular Location: Cell membrane
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: The tumor necrosis factor receptor (TNFR) superfamily represents a growing family of type I transmembrane glycoproteins that are involved in various cellular functions, including proliferation, differentiation and programmed cell death (1-3). These proteins share homology for cysteine-rich repeats in the extracellular ligand binding domain and an intracellular death domain (1-3). Members of the TNFR superfamily transmit signals through protein-protein interactions, and these signals can lead to the activation of either the caspase and Jun kinase pathways, which promote cell death, or the NFkB pathway, which results in cell survival (1). The ectodermal dysplasia receptor (EDAR) promotes all three of these pathways and mediates ectodermal differentiation (4). EDAR is encoded by the downless gene and is mutated in ectodermal dysplasia syndromes, which are characterized by impaired hair, teeth and sweat gland development (5). Ectodysplasin A (EDA) is a type II membrane protein that is encoded by the Tabby gene and produces many splice variants, the longest of which, EDA-A1, serves as the ligand for EDAR (5-7). EDA-A2, which differs from EDA-A1 by the deletion of two amino acids, binds only the X-linked ectodysplasin-A2 receptor (XEDAR) (7). Both EDAR and XEDAR exhibit homology with TROY (8).		

— VALIDATION IMAGES —

Tissue/cell: human gastric carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-EDA2R Polyclonal Antibody, Unconjugated(bs-7111R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

— SELECTED CITATIONS —

- **[IF=5.988]** Guo Zhijun. et al. Cisplatin-Induced Anorexia and Pica Behavior in Rats Enhanced by Chronic Stress Pretreatment. FRONT PHARMACOL. 2022 Jul;0:2565 WB ;Rat. 35910350
- **[IF=4.6]** Tokumasu Rino. et al. Transcription factor FoxO1 regulates myoepithelial cell diversity and growth. SCI REP-UK. 2024 Jan;14(1):1-15 IF, WB ;Mouse. 38212454