

bs-11094R**[Primary Antibody]****NCAM2 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit Clonality: Polyclonal GeneID: 4685 Target: NCAM2 Immunogen: KLH conjugated synthetic peptide derived from human NCAM2: 51-150/837. < Extracellular > Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: NCAM2 is an 837 amino acid protein encoded by the human gene NCAM2. NCAM2 contains five immunoglobulin-like domains, two Fibronectin type III domains, a transmembrane domain and a cytoplasmic domain. The gene is expressed most strongly in human adult and fetal brain. NCAM2 is a member of the neural cell adhesion molecule (NCAM) family. NCAMs are closely related cell surface glycoproteins involved in cell to cell interactions during growth and are thought to play an important role in embryogenesis and development. NCAM2 is considered a good candidate for involvement in certain Down syndrome phenotypes because a slight overexpression of NCAMs increases many-fold the homotypic adhesion properties of cells. Stat5 regulates NCAM2 in vivo by binding to the NCAM2 intron in the NKL natural killer cell line; this binding is induced by cytokines that activate Stat5. Neither Stat1 nor Stat3 bind to this region, despite sharing a consensus binding sequence with Stat5.	Isotype: IgG SWISS: O15394 Applications: Flow-Cyt (3ug/test) Reactivity: Human (predicted: Mouse, Rat, Rabbit) Predicted MW.: 91 kDa Subcellular Location: Cell membrane
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

— VALIDATION IMAGES —

Blank control: Hela. Primary Antibody (green line): Rabbit Anti-NCAM2 antibody (bs-11094R)

Dilution: 1µg /10⁶ cells; Isotype Control

Antibody (orange line): Rabbit IgG . Secondary

Antibody : Goat anti-rabbit IgG-PE Dilution: 3µg

/test. Protocol The cells were fixed with 4% PFA

(10min at room temperature)and then

permeabilized with 90% ice-cold methanol for

20 min at -20°C. The cells were then incubated in

5%BSA to block non-specific protein-protein

interactions for 30 min at at room temperature

.Cells stained with Primary Antibody for 30 min

at room temperature. The secondary antibody

used for 40 min at room temperature.

Acquisition of 20,000 events was performed.