bs-7949R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

AOC1 Rabbit pAb

GenelD: 26 **SWISS:** P19801

Target: AOC1

Immunogen: KLH conjugated synthetic peptide derived from human AOC1:

601-689/751.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Catalyzes the degradation of compounds such as putrescine,

histamine, spermine, and spermidine, substances involved in allergic and immune responses, cell proliferation, tissue

differentiation, tumor formation, and possibly apoptosis. Placental

DAO is thought to play a role in the regulation of the female

reproductive function.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500)

Reactivity: Mouse (predicted: Human,

Rat, Rabbit, Sheep, Cow,

Dog, Horse)

Predicted MW.: 83 kDa

Subcellular Secreted ,Extracellular

Location: matrix

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DAO1) Polyclonal Antibody, Unconjugated (bs-7949R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

— SELECTED CITATIONS —

• [IF=10.754] Apolloni S et al. Histaminergic transmission slows progression of amyotrophic lateral sclerosis. J Cachexia Sarcopenia Muscle. 2019 Apr 24. WB,ICC; Mouse. 31020811